Attach this document to your Epilepsy Management Plan if Midazolam is prescribed. This Emergency Plan should be completed by the prescribing doctor in consultation with the person and/or their family or carer. It must be attached to their epilepsy management plan which has been signed by their doctor. The Epilepsy Foundation recommends this plan be reviewed and signed by the person's doctor annually.





Emergency Medication Management Plan Midazolam (only to be administered by a trained person)

Midazolam Management Plan for (name)	
Date:	Date of birth:
Administration method: BUCCAL INTRANAS	SAL Nasal atomiser or Drip
1. FIRST DOSE Midazolam	
First dose = mg ml	
For single seizures:	
As soon as a	(seizure type) begins
☐ If the	(seizure type) continues longer than mins
For clusters of seizures:	
When (number)	(seizure type) occur/s within mins hrs
Other (please specify):	
Special instructions:	
2. SECOND DOSE Midazolam	
Second dose = mg r	nl
☐ Not prescribed OR	
☐ If the	(seizure type) continues for another mins following the first dose
When another (number) following the first dose	(seizure type) occur/s within mins hrs
Other (please specify):	
Special instructions:	
3. Maximum number of Midazolam dose	s to be given in a 24-hour period
Maximum number:	
4. Dial 000 to call the ambulance:	
☐ Prior to administering Midazolam	
☐ If the seizure has not stopped	minutes after giving the Midazolam
Other (please specify):	minates after giving the midazolam

Prescribing doctor or specialist	
lame of doctor:	
Гelephone:	Date:
Signature	
	Insert jpeg here
Any special instructions e.g. storage of	medication, when on outings etc. or people to contact if emergency
medication is given.	medication, when on outings etc. or people to contact if emergency Relationship:
Any special instructions e.g. storage of medication is given. Name:	
Any special instructions e.g. storage of medication is given. Name: Telephone:	Relationship:
Any special instructions e.g. storage of medication is given. Name:	Relationship:

- · Keep out of reach of children
- Protect from light and store at room temperature (below 25° C)

5. Describe what to do after Midazolam has been administered:

Regularly check the expiry date.

